

09/679,043
updated Search
L/cook 12/27/05.

d his

(FILE 'HOME' ENTERED AT 09:32:52 ON 27 DEC 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
09:33:17 ON 27 DEC 2005

L1 174 S TCII AND COBALAMIN?
L2 1 S L1 AND UNBOUND?
L3 28 S L1 AND FREE?
L4 13 DUPLICATE REMOVE L3 (15 DUPLICATES REMOVED)
L5 18 S (APO TCII)
L6 5 DUPLICATE REMOVE L5 (13 DUPLICATES REMOVED)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
10:01:24 ON 27 DEC 2005

L7 26 S (HOLO TCII)
L8 12 DUPLICATE REMOVE L7 (14 DUPLICATES REMOVED)
L9 4 S L8 AND APO?

=>



My List - 0 Help

[Main Search](#) | [Advanced Keyword Search](#) | [Search History](#)

Search: [Refine Search](#)

> You're searching: Scientific and Technical Information Center

Item Information

Subscriptions

Holdings

Browse Catalog

by title:

- Journal of clinical ...

MARC Display

Journal of clinical pathology.

Author: British Medical Association.
Association of Clinical Pathologists.

Imprint: London, British Medical Association. 1947-

URL: <http://proquest.umi.com/pqdweb?RQT=318&VName=PQD&clientid=19649&pmid=28111>
Click here to see full text or page images (9/1/96-) available via Proquest Direct.

Notes: "Available on ADONIS, v. 44, no. 1 (1991)-v. 55, no. 9 (2002)
"The Journal of the Association of Clinical Pathologists."
Preliminary number dated July 1945.

ISSN: 0021-9746

Subjects: Pathology -- Periodicals

Description: v. : ill.; 25 cm.

[Add to my list](#)

Subscription Summary

US Patent & Trademark Office

Location: US Patent & Trademark Office

Collection: Biotechnology and Chemical Library

Call No.: RB1 .J55

Copy No.: 1

Status: Not Currently Received

Media Type: SERIAL

Notes: Cancelled with Dec. 1999.

Main run: Vol. 36 No. 2 (Feb 1983) - Vol. 47 No. 12 (Dec 1993)
Vol. 48 No. 1 (Jan 1995) - Vol. 49 No. 11 (Nov 1996)
Vol. 50 No. 12 (Dec 1996) - Vol. 50 No. 1 (Jan 1997)
Vol. 50 No. 3 (Mar 1997) - Vol. 50 No. 5 (May 1997)
Vol. 50 No. 7 (Jul 1997) - Vol. 50 No. 8 (Aug 1997)
Vol. 50 No. 4 (Aug 1997)
Vol. 50 No. 9 (Sep 1997) - Vol. 50 No. 11 (Nov 1997)
Vol. 51 No. 1 (Jan 1998) - Vol. 51 No. 8 (Aug 1998)
Vol. 51 No. 10 (Oct 1998) - Vol. 51 No. 12 (Dec 1998)
Vol. 52 No. 1 (Jan 1999) - Vol. 52 No. 12 (Dec 1999)

Show all
items

US Patent & Trademark Office

Location: US Patent & Trademark Office

Collection: Biotechnology and Chemical Library Microfilm

Call No.: RB1 .J55 Microfilm

Copy No.: 1

Status: Not Currently Received

Media
Type: film

Notes: Cancelled with Dec. 1999.

Microfilm: v. 33 (1980) - v. 52 (1999)

Show all
items

Email: pamela.hoeft@uspto.gov to ask questions or make suggestions.

Horizon Information Portal 3.05

Brought to you by *Scientific and Technical Information Center*

ANSWER 2 OF 4 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN

AN 1996:278666 BIOSIS

DN PREV199699001022

TI Characterization of monoclonal antibodies to epitopes of human transcobalamin II.

AU Quadros, Edward V. [Reprint author]; Rothenberg, Sheldon P.; McLoughlin, Patricia

CS Dep. Med., Div. Hematol./Oncol., SUNY-Health Sci. Cent., Brooklyn, NY 11203, USA

SO Biochemical and Biophysical Research Communications, (1996) Vol. 222, No. 1, pp. 149-154.
CODEN: BBRCA9. ISSN: 0006-291X.

DT Article

LA English

ED Entered STN: 25 Jun 1996
Last Updated on STN: 15 Aug 1996

AB Cellular uptake of cobalamin (Cbl) is mediated by transcobalamin II (TCII), a Cbl binding protein in the plasma. The TCII-Cbl complex binds to a cell surface receptor and is internalized by endocytosis. We have generated monoclonal antibodies (mAbs) to human TCII that can be distinguished into three functional types on the basis of interaction with three different regions of the protein. Type 1: Receptor blocking. This mAb binds **holo-TCII** and inhibits the cellular uptake of Cbl. Type 2: Cbl blocking. This mAb binds **apo-TCII** at or near the Cbl binding domain and inhibits the formation of **holo-TCII**. Type 3: Precipitating. This mAb binds both **holo-TCII** and **apo-TCII** but does not interfere with Chi binding. Whereas type 1 and type 2 mAb, following incubation with TCII-(57Co)Cbl or **apo-TCII**, respectively, inhibit the uptake of radio-labeled Cbl by K562 cells, type 3 mAb has no such activity with either form of TCII. These properties of type 1 and type 2 mAb that inhibit the cellular uptake of Cbl, may serve to induce rapid Cbl deficiency and provide a model to study the effect of selective Cbl depletion on cell division and differentiation as well as on the pathways dependent on the two Cbl cofactors, methyl-Cbl and 5'-deoxyadenosyl-Cbl.

CC Cytology - Human 02508
Biochemistry studies - Vitamins 10063
Biochemistry studies - Proteins, peptides and amino acids 10064
Biochemistry studies - Carbohydrates 10068
Biophysics - Membrane phenomena 10508
Blood - Blood and lymph studies 15002
Pharmacology - General 22002
Immunology - General and methods 34502

IT Major Concepts
Biochemistry and Molecular Biophysics; Blood and Lymphatics (Transport and Circulation); Cell Biology; Immune System (Chemical Coordination and Homeostasis); Membranes (Cell Biology); Pharmacology

IT Chemicals & Biochemicals
TRANSCOBALAMIN II; METHYL-COBALAMIN; 5'-DEOXYADENOSYL-COBALAMIN

IT Miscellaneous Descriptors
CELL DIFFERENTIATION; CELL DIVISION; DRUG DESIGN; HUMAN K562 CELL; METHYL-COBALAMIN; TRANSCOBALAMIN II RECEPTOR BLOCKER; 5'-DEOXYADENOSYL-COBALAMIN

ORGN Classifier
Hominidae 86215
Super Taxa
Primates; Mammalia; Vertebrata; Chordata; Animalia
Organism Name
Hominidae
Taxa Notes
Animals, Chordates, Humans, Mammals, Primates, Vertebrates

RN 12651-28-4 (TRANSCOBALAMIN II)
13422-55-4 (METHYL-COBALAMIN)
13870-90-1 (5'-DEOXYADENOSYL-COBALAMIN)

ANSWER 2 OF 4 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN

AN 1996:278666 BIOSIS

DN PREV199699001022

TI Characterization of monoclonal antibodies to epitopes of human transcobalamin II.

AU Quadros, Edward V. [Reprint author]; Rothenberg, Sheldon P.; McLoughlin, Patricia

CS Dep. Med., Div. Hematol./Oncol., SUNY-Health Sci. Cent., Brooklyn, NY 11203, USA

SO Biochemical and Biophysical Research Communications, (1996) Vol. 222, No. 1, pp. 149-154.
CODEN: BBRCA9. ISSN: 0006-291X.

DT Article

LA English

ED Entered STN: 25 Jun 1996
Last Updated on STN: 15 Aug 1996

AB Cellular uptake of cobalamin (Cbl) is mediated by transcobalamin II (TCII), a Cbl binding protein in the plasma. The TCII-Cbl complex binds to a cell surface receptor and is internalized by endocytosis. We have generated monoclonal antibodies (mAbs) to human TCII that can be distinguished into three functional types on the basis of interaction with three different regions of the protein. Type 1: Receptor blocking. This mAb binds **holo-TCII** and inhibits the cellular uptake of Cbl. Type 2: Cbl blocking. This mAb binds **apo-TCII** at or near the Cbl binding domain and inhibits the formation of **holo-TCII**. Type 3: Precipitating. This mAb binds both **holo-TCII** and **apo-TCII** but does not interfere with Chi binding. Whereas type 1 and type 2 mAb, following incubation with TCII-(57Co)Cbl or **apo-TCII**, respectively, inhibit the uptake of radio-labeled Cbl by K562 cells, type 3 mAb has no such activity with either form of TCII. These properties of type 1 and type 2 mAb that inhibit the cellular uptake of Cbl, may serve to induce rapid Cbl depletion on cell division and differentiation as well as on the pathways dependent on the two Cbl cofactors, methyl-Cbl and 5'-deoxyadenosyl-Cbl.

CC Cytology - Human 02508
Biochemistry studies - Vitamins 10063
Biochemistry studies - Proteins, peptides and amino acids 10064
Biochemistry studies - Carbohydrates 10068
Biophysics - Membrane phenomena 10508
Blood - Blood and lymph studies 15002
Pharmacology - General 22002
Immunology - General and methods 34502

IT Major Concepts
Biochemistry and Molecular Biophysics; Blood and Lymphatics (Transport and Circulation); Cell Biology; Immune System (Chemical Coordination and Homeostasis); Membranes (Cell Biology); Pharmacology

IT Chemicals & Biochemicals
TRANSCOBALAMIN II; METHYL-COBALAMIN; 5'-DEOXYADENOSYL-COBALAMIN

IT Miscellaneous Descriptors
CELL DIFFERENTIATION; CELL DIVISION; DRUG DESIGN; HUMAN K562 CELL; METHYL-COBALAMIN; TRANSCOBALAMIN II RECEPTOR BLOCKER; 5'-DEOXYADENOSYL-COBALAMIN

ORGN Classifier
Hominidae 86215
Super Taxa
Primates; Mammalia; Vertebrata; Chordata; Animalia
Organism Name
Hominidae
Taxa Notes
Animals, Chordates, Humans, Mammals, Primates, Vertebrates

RN 12651-28-4 (TRANSCOBALAMIN II)
13422-55-4 (METHYL-COBALAMIN)
13870-90-1 (5'-DEOXYADENOSYL-COBALAMIN)

d his

(FILE 'HOME' ENTERED AT 09:32:52 ON 27 DEC 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
09:33:17 ON 27 DEC 2005

L1 174 S TCII AND COBALAMIN?
L2 1 S L1 AND UNBOUND?
L3 28 S L1 AND FREE?
L4 13 DUPLICATE REMOVE L3 (15 DUPLICATES REMOVED)
L5 18 S (APO TCII)
L6 5 DUPLICATE REMOVE L5 (13 DUPLICATES REMOVED)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
10:01:24 ON 27 DEC 2005

L7 26 S (HOLO TCII)
L8 12 DUPLICATE REMOVE L7 (14 DUPLICATES REMOVED)
L9 4 S L8 AND APO?

=>

d his

(FILE 'HOME' ENTERED AT 09:32:52 ON 27 DEC 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
09:33:17 ON 27 DEC 2005

L1 174 S TCII AND COBALAMIN?
L2 1 S L1 AND UNBOUND?
L3 28 S L1 AND FREE?
L4 13 DUPLICATE REMOVE L3 (15 DUPLICATES REMOVED)
L5 18 S (APO TCII)
L6 5 DUPLICATE REMOVE L5 (13 DUPLICATES REMOVED)

=>

d his

(FILE 'HOME' ENTERED AT 09:32:52 ON 27 DEC 2005)

FILE 'BIOSIS, CAPLUS, EMBASE, MEDLINE, CANCERLIT, JAPIO' ENTERED AT
09:33:17 ON 27 DEC 2005

L1	174 S TCII AND COBALAMIN?
L2	1 S L1 AND UNBOUND?
L3	28 S L1 AND FREE?
L4	13 DUPLICATE REMOVE L3 (15 DUPLICATES REMOVED)
L5	18 S (APO TCII)
L6	5 DUPLICATE REMOVE L5 (13 DUPLICATES REMOVED)

=>

ANSWER 3 OF 5 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN
 DUPLICATE 2
 AN 1996:278666 BIOSIS
 DN PREV199699001022
 TI Characterization of monoclonal antibodies to epitopes of human
 transcobalamin II.
 AU Quadros, Edward V. [Reprint author]; Rothenberg, Sheldon P.; McLoughlin,
 Patricia
 CS Dep. Med., Div. Hematol./Oncol., SUNY-Health Sci. Cent., Brooklyn, NY
 11203, USA
 SO Biochemical and Biophysical Research Communications, (1996) Vol. 222, No.
 1, pp. 149-154.
 CODEN: BBRC99. ISSN: 0006-291X.
 DT Article
 LA English
 ED Entered STN: 25 Jun 1996
 Last Updated on STN: 15 Aug 1996
 AB Cellular uptake of cobalamin (Cbl) is mediated by transcobalamin II
 (TCII), a Cbl binding protein in the plasma. The TCII-Cbl complex binds
 to a cell surface receptor and is internalized by endocytosis. We have
 generated monoclonal antibodies (mAbs) to human TCII that can be
 distinguished into three functional types on the basis of interaction with
 three different regions of the protein. Type 1: Receptor blocking. This
 mAb binds holo-TCII and inhibits the cellular uptake of Cbl. Type 2: Cbl
 blocking. This mAb binds **apo-TCII** at or near the Cbl
 binding domain and inhibits the formation of holo-TCII. Type 3:
 Precipitating. This mAb binds both holo-TCII and **apo-**
TCII but does not interfere with Chi binding. Whereas type 1 and
 type 2 mAb, following incubation with TCII-(57Co)Cbl or **apo-**
TCII, respectively, inhibit the uptake of radio-labeled Cbl by
 K562 cells, type 3 mAb has no such activity with either form of TCII.
 These properties of type 1 and type 2 mAb that inhibit the cellular uptake
 of Cbl, may serve to induce rapid Cbl deficiency and provide a model to
 study the effect of selective Cbl depletion on cell division and
 differentiation as well as on the pathways dependent on the two Cbl
 cofactors, methyl-Cbl and 5'-deoxyadenosyl-Cbl.
 CC Cytology - Human 02508
 Biochemistry studies - Vitamins 10063
 Biochemistry studies - Proteins, peptides and amino acids 10064
 Biochemistry studies - Carbohydrates 10068
 Biophysics - Membrane phenomena 10508
 Blood - Blood and lymph studies 15002
 Pharmacology - General 22002
 Immunology - General and methods 34502
 IT Major Concepts
 Biochemistry and Molecular Biophysics; Blood and Lymphatics (Transport
 and Circulation); Cell Biology; Immune System (Chemical Coordination
 and Homeostasis); Membranes (Cell Biology); Pharmacology
 IT Chemicals & Biochemicals
 TRANSCOBALAMIN II; METHYL-COBALAMIN; 5'-DEOXYADENOSYL-COBALAMIN
 IT Miscellaneous Descriptors
 CELL DIFFERENTIATION; CELL DIVISION; DRUG DESIGN; HUMAN K562 CELL;
 METHYL-COBALAMIN; TRANSCOBALAMIN II RECEPTOR BLOCKER;
 5'-DEOXYADENOSYL-COBALAMIN
 ORGN Classifier
 Hominidae 86215
 Super Taxa
 Primates; Mammalia; Vertebrata; Chordata; Animalia
 Organism Name
 Hominidae
 Taxa Notes
 Animals, Chordates, Humans, Mammals, Primates, Vertebrates
 RN 12651-28-4 (TRANSCOBALAMIN II)
 13422-55-4 (METHYL-COBALAMIN)

✓
 pull
 12/27/05

13870-90-1 (5'-DEOXYADENOSYL-COBALAMIN)

ANSWER 3 OF 5 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN

DUPLICATE 2

AN 1996:278666 BIOSIS

DN PREV199699001022

TI Characterization of monoclonal antibodies to epitopes of human transcobalamin II.

AU Quadros, Edward V. [Reprint author]; Rothenberg, Sheldon P.; McLoughlin, Patricia

CS Dep. Med., Div. Hematol./Oncol., SUNY-Health Sci. Cent., Brooklyn, NY 11203, USA

SO Biochemical and Biophysical Research Communications, (1996) Vol. 222, No. 1, pp. 149-154.

CODEN: BBRCA9. ISSN: 0006-291X.

DT Article

LA English

ED Entered STN: 25 Jun 1996

Last Updated on STN: 15 Aug 1996

AB Cellular uptake of cobalamin (Cbl) is mediated by transcobalamin II (TCII), a Cbl binding protein in the plasma. The TCII-Cbl complex binds to a cell surface receptor and is internalized by endocytosis. We have generated monoclonal antibodies (mAbs) to human TCII that can be distinguished into three functional types on the basis of interaction with three different regions of the protein. Type 1: Receptor blocking. This mAb binds holo-TCII and inhibits the cellular uptake of Cbl. Type 2: Cbl blocking. This mAb binds apo-TCII at or near the Cbl binding domain and inhibits the formation of holo-TCII. Type 3: Precipitating. This mAb binds both holo-TCII and apo-TCII but does not interfere with Chi binding. Whereas type 1 and type 2 mAb, following incubation with TCII-(57Co)Cbl or apo-TCII, respectively, inhibit the uptake of radio-labeled Cbl by K562 cells, type 3 mAb has no such activity with either form of TCII. These properties of type 1 and type 2 mAb that inhibit the cellular uptake of Cbl, may serve to induce rapid Cbl deficiency and provide a model to study the effect of selective Cbl depletion on cell division and differentiation as well as on the pathways dependent on the two Cbl cofactors, methyl-Cbl and 5'-deoxyadenosyl-Cbl.

CC Cytology - Human 02508

Biochemistry studies - Vitamins 10063

Biochemistry studies - Proteins, peptides and amino acids 10064

Biochemistry studies - Carbohydrates 10068

Biophysics - Membrane phenomena 10508

Blood - Blood and lymph studies 15002

Pharmacology - General 22002

Immunology - General and methods 34502

IT Major Concepts

Biochemistry and Molecular Biophysics; Blood and Lymphatics (Transport and Circulation); Cell Biology; Immune System (Chemical Coordination and Homeostasis); Membranes (Cell Biology); Pharmacology

IT Chemicals & Biochemicals

TRANSCOBALAMIN II; METHYL-COBALAMIN; 5'-DEOXYADENOSYL-COBALAMIN

IT Miscellaneous Descriptors

CELL DIFFERENTIATION; CELL DIVISION; DRUG DESIGN; HUMAN K562 CELL; METHYL-COBALAMIN; TRANSCOBALAMIN II RECEPTOR BLOCKER; 5'-DEOXYADENOSYL-COBALAMIN

ORGN Classifier

Hominidae 86215

Super Taxa

Primates; Mammalia; Vertebrata; Chordata; Animalia

Organism Name

Hominidae

Taxa Notes

Animals, Chordates, Humans, Mammals, Primates, Vertebrates

RN 12651-28-4 (TRANSCOBALAMIN II)

13422-55-4 (METHYL-COBALAMIN)

13870-90-1 (5'-DEOXYADENOSYL-COBALAMIN)

ANSWER 4 OF 5 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN
DUPLICATE 3

AN 1984:271994 BIOSIS
DN PREV198478008474; BA78:8474
TI LIMITATIONS OF PROTEIN COATED CHARCOAL IN THE SEPARATION OF FREE FROM
BOUND VITAMIN B-12 IN SERUM.
AU JACOB E [Reprint author]; WONG K-T J
CS DEP MED, ST BONIFACE HOSP, 409 TACHE AVE, WINNIPEG, MANITOBA R2H 2A6, CAN
SO Journal of Clinical Pathology (London), (1983) Vol. 36, No. 9, pp.
1022-1027.
CODEN: JCPAAK. ISSN: 0021-9746.
DT Article
FS BA
LA ENGLISH
AB The effect of Hb and albumin-coated charcoal on the concentration of
vitamin B12 binding proteins in [human] serum was investigated. As
commonly employed, coated charcoal removes a significant amount of
transcobalamin II (TCII) from serum, but does not affect transcobalamin I
and III (TCI and III). Increasing the protein coat up to about 10 mg of
protein/25 mg of charcoal reduces the adsorption of TCII, but increasing
the protein concentration beyond this has little added effect. The amount
of adsorption is proportional to the amount of coated charcoal employed,
but even small amounts adsorb some TCII. Protein-coated charcoal is not
the ideal way of separating free from bound vitamin B12 in serum; it
cannot reliably be used for the measurement of the concentration of
apo-TCII but can be employed for the measurement of
apo-TCI and III.
CC Clinical biochemistry - General methods and applications 10006
Biochemistry methods - Vitamins 10053
Biochemistry methods - Proteins, peptides and amino acids 10054
Biochemistry studies - Vitamins 10063
Biochemistry studies - Proteins, peptides and amino acids 10064
Biochemistry studies - Porphyrins and bile pigments 10065
Biophysics - Methods and techniques 10504
Movement 12100
Blood - General and methods 15001
Blood - Blood cell studies 15004
IT Major Concepts
Biochemistry and Molecular Biophysics; Blood and Lymphatics (Transport
and Circulation); Clinical Chemistry (Allied Medical Sciences)
IT Miscellaneous Descriptors
HUMAN TRANS COBALAMINS/
ORGN Classifier
Hominidae 86215
Super Taxa
Primates; Mammalia; Vertebrata; Chordata; Animalia
Taxa Notes
Animals, Chordates, Humans, Mammals, Primates, Vertebrates
RN 68-19-9 (VITAMIN B12)
12774-24-2D (TRANSCOBALAMINS)

ANSWER 4 OF 5 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN
DUPLICATE 3

AN 1984:271994 BIOSIS

DN PREV198478008474; BA78:8474

TI LIMITATIONS OF PROTEIN COATED CHARCOAL IN THE SEPARATION OF FREE FROM
BOUND VITAMIN B-12 IN SERUM.

AU JACOB E [Reprint author]; WONG K-T J

CS DEP MED, ST BONIFACE HOSP, 409 TACHE AVE, WINNIPEG, MANITOBA R2H 2A6, CAN

SO Journal of Clinical Pathology (London), (1983) Vol. 36, No. 9, pp.
1022-1027.

CODEN: JCPAAK. ISSN: 0021-9746.

DT Article

FS BA

LA ENGLISH

AB The effect of Hb and albumin-coated charcoal on the concentration of
vitamin B12 binding proteins in [human] serum was investigated. As
commonly employed, coated charcoal removes a significant amount of
transcobalamin II (TCII) from serum, but does not affect transcobalamin I
and III (TCI and III). Increasing the protein coat up to about 10 mg of
protein/25 mg of charcoal reduces the adsorption of TCII, but increasing
the protein concentration beyond this has little added effect. The amount
of adsorption is proportional to the amount of coated charcoal employed,
but even small amounts adsorb some TCII. Protein-coated charcoal is not
the ideal way of separating free from bound vitamin B12 in serum; it
cannot reliably be used for the measurement of the concentration of
apo-TCII but can be employed for the measurement of
apo-TCI and III.

CC Clinical biochemistry - General methods and applications 10006

Biochemistry methods - Vitamins 10053

Biochemistry methods - Proteins, peptides and amino acids 10054

Biochemistry studies - Vitamins 10063

Biochemistry studies - Proteins, peptides and amino acids 10064

Biochemistry studies - Porphyrins and bile pigments 10065

Biophysics - Methods and techniques 10504

Movement 12100

Blood - General and methods 15001

Blood - Blood cell studies 15004

IT Major Concepts

Biochemistry and Molecular Biophysics; Blood and Lymphatics (Transport
and Circulation); Clinical Chemistry (Allied Medical Sciences)

IT Miscellaneous Descriptors

HUMAN TRANS COBALAMINS/

ORGN Classifier

Hominidae 86215

Super Taxa

Primates; Mammalia; Vertebrata; Chordata; Animalia

Taxa Notes

Animals, Chordates, Humans, Mammals, Primates, Vertebrates

RN 68-19-9 (VITAMIN B12)

12774-24-2D (TRANSCOBALAMINS)